

B4: Portions of this document are withheld due to copyright laws.

## Metabolites of a Tobacco-specific Nitrosamine, 4-(Methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK), in the Urine of Smokeless Tobacco Users: Relationship between Urinary Biomarkers and Oral Leukoplakia<sup>1</sup>

Laura A. Kresty, Steven G. Carmella, Anna Borukhova, Shobha A. Akerkar, Rajaram Gopalakrishnan, Randall E. Harris, Gary D. Stoner, and Stephen S. Hecht<sup>2</sup>

Department of Preventive Medicine, Ohio State University, Columbus, Ohio 43210-1240 [L. A. K., R. G., R. E. H., G. D. S.], and American Health Foundation, Valhalla, New York 10595 [S. G. C., A. B., S. A. A., S. S. H.]

(b) (4)

(b) (4)

Received 12/4/95; revised 3/19/96; accepted 3/26/96.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>1</sup> Supported by National Cancer Institute Grant CA-46535. Cotinine and creatinine analyses were carried out in the American Health Foundation Clinical Biochemistry Facility, and statistical analyses were performed in part by the Biostatistics and Computer Facility, supported in part by National Cancer Institute Cancer Center Support Grant CA-17613.

<sup>2</sup> To whom requests for reprints should be addressed, at University of Minnesota Cancer Center, Box 806 UMHC, 420 Delaware St., S.E., Minneapolis, MN 55455.