

Smokeless tobacco use in Sweden and other 17 European countries

Maria E. Leon¹, Alessandra Lugo², Paolo Boffetta³, Anna Gilmore⁴, Hana Ross⁵, Joachim Schüz¹, Carlo La Vecchia⁶, Silvano Gallus²

- 1 Section of Environment and Radiation, International Agency for Research on Cancer (IARC), Lyon, France
- 2 Department of Epidemiology, IRCCS—Istituto di Ricerche Farmacologiche “Mario Negri”, Milan, Italy
- 3 Icahn School of Medicine at Mount Sinai, Tisch Cancer Institute and Institute for Translational Epidemiology, New York, USA
- 4 Department for Health and UK Centre for Tobacco and Alcohol Studies, University of Bath, Bath, UK
- 5 School of Economics, University of Cape Town, Cape Town, South Africa
- 6 Department of Clinical Sciences and Community Health, Università degli Studi di Milano, Milan, Italy

Correspondence: Maria E. Leon, Section of Environment and Radiation, International Agency for Research on Cancer (IARC), 150 Cours Albert Thomas, Lyon 69008, France, Tel: +33-4-72738171, Fax: +33-4-72738320, e-mail: leonrouxm@iarc.fr

(b) (4)



(b) (4)

