

Evidence That Smokeless Tobacco Use Is a Gateway for Smoking Initiation in Young Adult Males¹

C. Keith Haddock, Ph.D.,*² Mark Vander Weg, Ph.D.,† Margaret DeBon, Ph.D.,† Robert C. Klesges, Ph.D.,† G. Wayne Talcott, Ph.D.,‡ Harry Lando, Ph.D.,§ and Alan Peterson, Ph.D.||

**University of Missouri, Kansas City, and Mid America Heart Institute, St. Luke's Hospital, Kansas City, Missouri 64110;*

†University of Memphis, Memphis, Tennessee 38119; ‡Office of the Air Force Surgeon General, Bolling Air Force Base, Maryland 20332;

§University of Minnesota, Minneapolis, Minnesota 55454; and Wilford Hall Medical Center, Lackland Air Force Base, Texas 78236

(b) (4)



(b) (4)



¹ This paper was supported by a grant from the U.S. National Institutes of Health (HL-53478) awarded to the University of Memphis (R.C. Klesges, Principal Investigator). The views expressed in the article are those of the authors and are not the official policy of the U.S. Government, Department of Defense, or U.S. Air Force.

² To whom requests for reprints should be addressed at Health Research Group, University of Missouri, Kansas City, 4825 Troost, Suite 124, Kansas City, MO 64110.

